

Identification of the *Propionibacterium acnes* using Polymerase Chain Reaction (PCR) in the Acne vulgaris lesions

Negin Naghdi^{1*}, Masood Ghane¹

1. Department of Microbiology, Faculty of Biological Sciences, Islamic Azad University, Tonekabon Branch, Tonekabon, Iran

Abstract

Background: Acne vulgaris is an inflammatory chronic disease of pilosebaceous unit. One of the most important factors playing a role in occurrence of acne is presence of *Propionibacterium acnes*. With the aim of molecular identification of the *P. acnes* from the acne vulgaris lesions, current research was carried out.

Methods: In this study, contents within the lesions was collected from 70 patients. The presence of the *P. acnes* was examined by a specific PCR technique.

Results: Of 70 samples, 58 samples (82.85%) were determined to be positive in terms of presence of *P. acnes*. No significant relationship was observed between presence of *P. acnes* and each one of the studied demographic factors, including gender, age, disease period, family background and treatment background.

Conclusions: The adopted molecular technique has obviated the limitations associated with the culture method for identification of the bacteria. To overcome the problems with conventional culture techniques for *P. acnes*, this PCR method is promising for better identification of this bacterium.

Keywords: Identification, Polymerase Chain Reaction (PCR), *Propionibacterium acnes*, Acne vulgaris

Introduction

Acne vulgaris is the most prevalent skin disorder of follicle sebaceous (Bhambri et al., 2009) that observed in 80% of the juveniles (Ray et al., 2013; Jahns et al., 2012) at the age ranged from 14 to 19 years old. Due to earlier onset of the puberty age, this disease is more prevalent in the women, while it's more acute shapes are observed in the man (Adityan and

Thappa, 2009). Pathogenesis of acne is multifactorial; obstruction of sebaceous pores, increase of producing the sebum, *P.acnes*, inflammation and androgens are the factors supposed to play a role in formation of the acne (Bhambri et al., 2009). For more than a century, the clinical relationship between *P. acnes* and appearance of the acne vulgaris has been identified (McDowell et al., 2013). So that increase of number of this bacterium leads to acuteness of the acne and decrease of number of that leads to improvement of the

***Corresponding author. Negin Naghdi, MSc**
Department of Microbiology, Faculty of Biological sciences,
Islamic Azad University, Tonekabon Branch, Tonekabon, Iran
Tel. +9891129220891
Email: negin.naghdi91@yahoo.com

acne (Bhambri, et al., 2009).

P. acnes is a polymorphic (Cauich et al., 2001; Perry and Lambert, 2006), gram positive, facultative anaerobic, without spore (Cauich, et al., 2001; Skinner et al., 1978), non motile (Skinner et al., 1978) and slowly growing bacilli (Portillo et al., 2013; Perry and Lambert, 2006). It has the best growth in the oxygen limitation conditions (Cove et al., 1983) and rate of its growth decreases in the high concentration of oxygen (Perry and Lambert, 2006; Cove et al., 1983). This bacterium is an opportunistic pathogen (McDowell et al., 2013) and, through extracellular products, including Hyaluronidase, Protease and Lipase and, also, Chemotaxis factors for the Neutrophils, Lymphocytes and Macrophages, plays a role in the pathogenesis of acne (Burkhart et al., 1999).

Today, PCR technique is used to diagnose an extensive spectrum of the biological factors. Compared to the traditional methods of diagnosis, PCR method is of a high speed, accuracy and specificity and sensitivity. Considering the weak points of the traditional methods in this study, PCR method was used in order to diagnose quickly and accurately *P.acnes* in the lesions isolated from the patients with acne vulgaris.

Materials and Methods

Sample collection

The current research has been carried out cross

sectional analysis. In this study, 70 patients with acne vulgaris referred to the skin clinic in the Tonekabon township within the years ranged from 2013 to 2014 were studied. Of total 70 collected samples, 49 cases were female and 21 cases were male who placed in the age limits ranged from 14 to 32 years. The studied samples were collected from the face (forehead, cheek and chin) of the studied individuals. The collected samples were transferred to the micro tube containing TBE buffer (Tris-borate-EDTA buffer) and kept under -20°C until enforcement of the DNA extraction process. Of a number of the samples, microbic culture was prepared randomly. After identification of bacillus of *P.acne*, the purified colonies were used to perform process of sequencing.

DNA extraction

300 µl of the taken collected samples were solved in 200 µl of the TE buffer (Tris-EDTA buffer) and, following the addition of 2 µl of lysozyme, kept for 30 minutes in the room temperature. Then, DNA extraction was conducted by use of the instruction of maker company of the extraction kit (Qiagen, Germany). In order to study accuracy of the DNA extraction, photoabsorption of the samples in the range of 260 and 280 nm was evaluated by biophotometer apparatus (Eppendorf, Germany).

PCR

PAR-1 and PAR-2 specific primer made by TAG Copenhagen company Denmark and PCR kit (Qiagen-Germany) were used to perform PCR. PCR mix with final volume of 20 μ l including, 10 μ l of Master Mix along with 0.5 μ l of Forward primer, 0.5 μ l of Reverse primer (10 pmol), 5 μ l of extracted DNA and 4 μ l of sterile distilled water was prepared in the 0.2

micro tube and placed inside the thermocycler apparatus (Eppendorf-Germany). PCR reaction was carried out in 35 cycles, including denaturation in 95°C for 60 seconds, primer's annealing in 60°C for 30 seconds and extension in 72°C for 90 seconds. Also, initial denaturation with 95°C for 5 minutes and final extension with 72°C for 10 minutes were taken into consideration (Niazi et al., 2010).

Table 1 Sequence of primer 5'→ 3'

Primer	Sequence of primer 5'→ 3'	PCR Product	References
PAR-1	5'-AGC TCG GTG GGG TTC TCT CAT C-3'	1202 bp	Niazi et al., 2010
PAR-2	5'-GCT TCC TCA TAC CAC TGG TCA TC-3'		

Electrophoresis

Electrophoresis was performed using 5 μ l of each samples. A DNA molecular ladder (1Kb) was used as control (Qiagen-Germany). The electrophoresis was done on 1.5% agarose gel adjusting voltage of 75 V for 40 min. Then, the gel was transferred to Transilluminator apparatus (Gel Doc-England) to visualize the DNA bands.

were selected for sequencing. 50 μ l of the PCR product obtained by specific primer of *P. acnes* were transferred to Korea (Macrogen Company, Korea) for sequencing and identification of the isolated strains. Then, the obtained gene sequence was analyzed using BLAST software available in NCBI site.

Sequencing

In order to confirm PCR data, four samples

Results

Of 70 samples studied by use of PCR technique, 58 positive samples (82.85%) were identified in terms of *P. acnes*.




Figure 1 Amplification products separated on agarose gel.

Electrophoresis was performed on 1.5% agarose gel. Lane 1: negative control, Lane 2: DNA molecular ladder (1kb), Lane 3: positive control and Lanes 4 to 7: *P. acne*-positive samples.

The purified colonies of the *P. acne* were used in order to sequencing. Figure 2 is the microscopic shape of the studied bacterium. The results achieved from the sequencing, also, confirmed result of PCR. In table 2, the results achieved from sequencing of the samples evaluated using BLAST software

available in NCBI site are observed.




Figure 2 Gram's staining showing Gram positive bacillus of *P. acne*

Table 2 Sequencing data of the samples observed in NCBI site

Sample	<i>P. acnes</i> Strain	Sequence ID
1	hdn-1	CP006032.1
2	PA44	KJ572677.1
3	hdn-1	CP006032.1
4	PA62	KJ572678.1

Findings from research on the basis of demographic information, including gender, age, and disease period, background of acne in family and background of treatment were analyzed. In order to interpret the results, SPSS

statistical software (SPSS version 18) and Chi-square independence test were used. The studies showed that there is not any relationship between rate of frequency of the *P.acnes* and each one of the studied factors.

Table 3 Demographic information of the studied patients and relative and absolute frequency of the *P.acnes* in terms of each one of the mentioned factors

Index	Status	Frequency	Percentage frequency (%)	Positive PCR	Percentage Positive PCR (%)	P-value
gender	Female	49	70	42	85.7	0.261
	Male	21	30	16	76.2	
Age (years)	10-15	6	8.57	6	100	0.330
	15-20	32	45.71	27	84.4	
	20-25	18	25.71	13	72.2	
	25-30	12	17.14	10	83.3	
	30-35	2	2.86	2	100	
Time Disease (years)	Random	37	52.86	33	89.2	0.398
	1-5	20	28.57	15	75	
	5-10	11	15.71	8	72.7	
	10-15	2	2.86	2	100	
Background of acne in family	Yes	37	52.86	31	83.8	0.828
	No	33	47.14	27	81.8	
Background of treatment	Yes	29	41.43	22	37.9	0.215
	No	41	58.57	36	58.3	

Discussion

In the recent years, our knowledge regarding microorganisms has been depended on the culture technique, while culture method is able to identify little minority (less than 1%) of the bacteria (Grice et al., 2008). For the purpose of identification of the viable but non culturable (VBNC), microorganisms which are not capable of growing in the usual culture media (Kong and Segre, 2012; Rhoads et al., 2012), microorganisms growing slowly (Millar et al., 2007) and, also, obligate anaerobes microorganisms (Rhoads et al., 2012) the molecular techniques have a lot of applications.

Usage of the techniques relied on nucleic acid has obviated the limitations available on the way of identification of microorganisms through culture technique (Taravati et al., 2013). In the molecular methods, little quantity of the microorganism's DNA is identifiable and living of the studied microorganism is not to be required (Rhoads et al., 2012; Kong and Segre, 2012). Also, in order to carry out PCR, samples such as blood and liquids of the body can be used, and preparation of the fresh sample is not required. Before sampling, antibiotic consumption doesn't influence on the result achieved from PCR, while it leads to establishment of the false negative response in the results obtained from culture.

P. acnes is a fastidious bacterium which requires enrich medium, special nutrient

factors and anaerobic conditions in order to grow. Due to its slow growth, identification of the *P. acnes* through culture is not so much possible. With aim of molecular identification of the *P. acnes* assisted with PCR method in the acne vulgaris lesions isolated from the patients with acne, the current study was accomplished. Of 70 studied samples, 58 positive samples (82.85%) were identified in terms of presence of the *P. acnes* with assistance of PCR technique, suggesting high outbreak of the bacterium in the acne lesions. In 2003, a research was conducted by Le Page et al aiming at diagnosis of the vascular prosthesis infection through method amplification and sequencing of *16S rDNA*. In this study, of 20 studied samples, 5 positive samples (25%) of the *P. acnes* were identified with assistance of the PCR method (Le Page et al., 2003). In 2006, a research was conducted on the individuals with endophthalmitis by Bagyalakshmi et al. 30 samples were collected from the studied patients and using mPCR method, 4 positive samples (13.3%) of the *P. acnes* were diagnosed (Bagyalakshmi et al., 2006). In 2013, a study was carried out by Rollason et al, on the patients with herniated waist disk surgery. Of total 64 studied samples, presence of the *P. acnes* in 24 samples (38%) was confirmed with aid of analysis of *recA* gene sequence (Rollason, et al., 2013). In 2014, a study was carried out by Bunker et al

on the individuals with shoulder infection following the orthopedic surgery. Of 10 studied patients, 6 positive samples (60%) of *P. acnes* were diagnosed by use of PCR method (Bunker et al., 2014).

Some discrepancies are observed among the results achieved from the researches, this discrepancy may be related to sampling place, type of the studied sample (acne or other infections resulted from *P. acnes*) and conditions of PCR performance. With a general glance at the results achieved from the current study and other conducted studies, it can be concluded that, using PCR method, higher percentage of the microorganisms is to be identified. Also, usage of the molecular methods for the purpose of identification of the biologic factors compared to the traditional methods of diagnosis leads to time and cost economizing. Despite the day increasing improvements of molecular techniques for the purpose of diagnosis of a variety of infections, culture method as a standard method remains to be a special value and position (Taravati et al., 2013; Kong and Segre, 2012).

Conclusion

With regard to the existed limitations for identification of microorganisms through culture technique as well as high speed and accuracy of the PCR technique in the diagnosis of pathogenic factors, this method, can be

considered an appropriate tool for identification of the studied bacterium.

References

- [1] Adityan B, Thappa DM. 2009. Profile of acne vulgaris-A hospital-based study from South India. *Indian Journal of Dermatology, Venereology and Leprology*, 75(3): 272-8.
- [2] Bagyalakshmi R, Madhavan HN, Therese Kl. 2006. Development and application of multiplex polymerase chain reaction for the etiological diagnosis of infectious endophthalmitis. *Journal of Postgraduate Medicine*, 52(3): 179-82.
- [3] Bhambri S, Del Rosso JQ, Bhambri A. 2009. Pathogenesis of acne vulgaris: recent advances. *J Drugs Dermatol*, 8(7): 615-8.
- [4] Bunker TD, Boyd M, Gallacher S, Auckland CR, Kitson J, Smith CD. 2014. Association between *Propionibacterium acnes* and frozen shoulder: a pilot study. *Shoulder and Elbow*, 6(4): 257-61.
- [5] Burkhart CG, Burkhart CN, Lehmann PF. 1999. Acne: A review of immunologic and microbiologic factors. *Postgraduate Medical Journal*, 75(884): 328-31.
- [6] Cauich P, Alatríste F, García E, Aquino C. 2001. Identification of anaerobic nonspore forming gram-positive bacilli by biochemical tests and gas-liquid chromatography. *Revista Latinoamericana de Microbiología*, 43(1): 27-35.

- [7] Cove JH, Holland KT, Cunliffe WJ. 1983. Effects of oxygen concentration on biomass production, maximum specific growth rate and extracellular enzyme production by three species of cutaneous propionibacteria grown in continuous culture. *Journal of General Microbiology*, 129 (11): 3327-34.
- [8] Grice EA, Kong HH, Renaud G, Young AC, Bouffard GG, Blakesley RW, Wolfsberg TG, Turner ML, Segre JA. 2008. A diversity profile of the human skin microbiota. *Genome Research*, 18(7): 1043-50.
- [9] Jahns AC, Lundskog B, Ganceviciene R, Palmer RH, Golovleva I, Zouboulis CC, McDowell A, Patrick S, Alexeyev OA. 2012. An increased incidence of *Propionibacterium acnes* biofilms in acne vulgaris: A case-control study. *British Journal of Dermatology*, 167(1): 50-8.
- [10] Kong HH, Segre JA. 2012. Skin Microbiome: Looking Back to Move Forward. *Journal of Investigative Dermatology*, 132(3): 933-9.
- [11] Le Page L, Podglajen I, Chemla E, Mainardi JL. 2003. Molecular diagnosis of a vascular prosthesis infection, due to *Propionibacterium acnes*, by amplification and sequencing of 16S rDNA. *Clinical Microbiology and Infection*, 9(11): 1125-7.
- [12] McDowell A, Patrick S, Eishi Y, Lambert P, Eady A. 2013. *Propionibacterium acnes* in human health and disease. *BioMed Research International*, 2013: 3 pages.
- [13] Millar BC, Xu J, Moore JE. 2007. Molecular diagnostics of medically important bacterial infections. *Molecular Diagnostics: Current Technology and Applications*, 9(1): 21-39.
- [14] Niazi SA, Clarke D, Do T, Gilbert SC, Mannocci F, Beighton D. 2010. *Propionibacterium acnes* and *Staphylococcus epidermidis* isolated from refractory endodontic lesions are opportunistic pathogens. *Journal of Clinical Microbiology*, 48(11): 3859-69.
- [15] Perry AL, Lambert PA. 2006. *Propionibacterium acnes*. *Letters in Applied Microbiology*, 42(3): 185-8.
- [16] Portillo ME, Corvec S, Borens O, Trampuz A. 2013. *Propionibacterium acnes*: an underestimated pathogen in implant-associated infections. *Bio Med Research International*, 2013: 10 pages.
- [17] Ray C, Trivedi P, Sharma V. 2013. Acne and its treatment. *International Journal of Research in Pharmaceutical and Biosciences*, 3(1): 1-16.
- [18] Rhoads DD, Wolcott RD, Sun Y, Dowd SE. 2012. Comparison of culture and molecular identification of bacteria in chronic wounds. *International Journal of Molecular Sciences*, 13(3): 2535-50.
- [19] Rollason J, McDowell A, Albert HB, Barnard E, Worthington T, Hilton AC,

- Vernallis A, Patrick S, Elliott T, Lambert P. 2013. Genotypic and antimicrobial characterisation of *Propionibacterium acnes* isolates from surgically excised lumbar disc herniations. *BioMed Research International*, 2013: 1-7.
- [20] Skinner PR, Taylor AJ, Coakham H. 1978. Propionibacteria as a cause of shunt and postneurosurgical infections. *Journal of Clinical Pathology*, 31(11): 1085-90.
- [21] Taravati P, Lam D, Van Gelder RN. 2013. Role of molecular diagnostics in ocular microbiology. *Current Ophthalmology Reports*, 1(4): 181-9.